	TA	ILGATE/TOOL	BOX SAFET	Y MEETINGS		
SAFETY S SERVICES COMPANY	afety Services Compan	y-Safety Meeting Division,	PO Box 78402, Coron	a, CA 92877 Toll Free (866	)204-4786	
			Job Site Location	Job Site Location:		
Date:	Start Tir	ne: Finish Time:	Foreman/Supe	rvisor:		
		Topic 21: First				
result in mi	ssed work and needless	mong the most common inju pain. A little care to prepar uscle strain can help you rec	ries suffered in the wo	rkplace. Sore backs, stiff no the number of injuries due t	o muscle strains. Proper	
most effecti for 3 to 10 r count of ten efficiently, a	ve method of warming minutes. A person shou . Stretching lengthens i	erson should stretch only aft up is slowly going through to all stretch after warming up muscles and tendons, and all order to avoid muscle strains the tips:	the actual motions of the or exercising, and each ows longer muscles to	he exercise or work to be pe 1 stretch should be comforta generate more force around	rformed at a relaxed pac ble enough to hold for a the joints, contract more	
$\square$	Stretch your hamstrir Stretch your quadrice Stretch your Achilles Stretch your arms and	eps tendons		Stretch your neck Stretch your groin Stretch your back		
ce, compression of the conclusion of the conclus	Immediately stop the minimize swelling an Ice causes the blood pack or chemical cold over a period of 1 to 1 Wrapping the injure part above the level o Do not place hot pack Place hot packs on the Anti inflammatory a health care profession Begin light stretchin If the muscle strain d A program of planned muscles. Strengthenin	e activity that caused the straid prevent the injury from bevessels to constrict, helping to pack on the affected area for 1½ hours. This process can be part with an elastic bandar of the heart (elevation) helps less on the affected area for at a faffected area for 15 to 20 m gents like aspirin or ibuprof al before taking any medicate of the affected area 72 hou oes not improve within 72 to physical exercise will help to g and stretching your muscle alt your physician before beg	tin. The injured part is coming worse. to limit inflammation a per 10 minutes, then remove repeated several time age (compression) and limit swelling. least 24 hours (heat within the every 2 hours are may reduce swelling tion). The after the injury. The per per prevent must be swell help pr	rested immediately to and reduce pain. Place an ic nove it for 10 minutes, repeates over the first 24 hours. raising the injured all increase swelling and sevofter the first 24 hours g and pain (seek the advice cian.	e at erity of the injury).	
pecific Wo	rk-Site Hazards and Sa	fety Suggestions:				
Employee	Signatures:	(My signature attests and veri regulations, and that I have t	ifies my understanding of a not suffered, experienced, o	nd agreement to comply with, all c r sustained any recent, reportable,	ompany safety policies and job-related injury or illness.)	